




Slide 1 of 2

| Metric       |            | CQC Key Question   | 2017 Report <sup>1</sup> | National Aggregate (England only) | National Standard | Comparison to other sites |   |
|--------------|------------|--|--------------------------|-----------------------------------|-------------------|---------------------------|---|
| Trust-level  | 5205 cases | Case ascertainment   | Well-Led                 | <b>101.01%*</b>                   | 92.00%            | >95%                      | Meets the national standard   |
| Ante-natal   | 344 cases  | Case-mix adjusted proportion of small-for-gestational-age babies (birthweight below 10th centile) who are not delivered before their due date                  | Effective                | <b>58.18%</b>                     | 55.30%            | N/A                       | <br>38.7   74.9<br>Within expected range |
|              |            | Case-mix adjusted proportion of elective deliveries (caesarean or induction) between 37 and 39 weeks with no documented clinical indication for early delivery | Effective                | <b>36.20%</b>                     | 26.00%            | N/A                       | <br>3.6   55.6<br>Within expected range  |
| Intra-partum | 4642 cases | Case-mix adjusted overall caesarean section rate for single, term babies   | Effective                | <b>20.16%</b>                     | 24.50%            | N/A                       | <br>20.2   32.4<br>Lower than expected   |



**KEY ONLY**

Lower than expected (above 99.8% control limit) | Higher than expected (above 99.8% control limit) | Lower than expected (below 99.8% control limit) | Higher than expected (above 99.8% control limit)

Trust | Trust

Within expected range | Within expected range

Display of performance boundaries does vary depending on volume of activity. See FAQs for further information.

Slide 2 of 2


|              | Metric  | CQC Key Question | 2017 Report <sup>1</sup> | National Aggregate (England only) | National Aspirational Standard | Comparison to other sites |
|--------------|---|------------------|--------------------------|-----------------------------------|--------------------------------|---------------------------|
| Intra-partum | 4579 cases<br>Case-mix adjusted proportion of single, term infants with a 5-minute Apgar score of less than 7               | Effective        | 1.21%*                   | 1.20%                             | N/A                            | Within expected range     |
|              | 3617 cases<br>Case-mix adjusted proportion of vaginal births with a 3rd/4th degree perineal tear                            | Safe             | 3.03%*                   | 3.60%                             | N/A                            | Within expected range     |
|              | 4642 cases<br>Case-mix adjusted proportion of women with severe post partum haemorrhage of greater than or equal to 1500 ml | Safe             | 3.50%*                   | 2.70%                             | N/A                            | Within expected range     |
| Post-partum  | 4997 cases<br>Proportion of live born babies who received breast milk for the first feed                                    | Effective        | 76.93%                   | 74.10%                            | N/A                            | Within expected range     |



Produced by HQIP in partnership with the



KEY ONLY



Display of performance boundaries does vary depending on volume of activity. See FAQs for further information.