

Metric	CQC Key Question	Oct 2018 and Mar 2019	National Aggregate (England & Wales)	National Aspirational Standard	Comparison to other hospitals	
125 Cases	Case ascertainment	Well-led	Not yet available	Not yet available		
125 Cases	Percentage of patients seen by a member of the respiratory team within 24hrs of admission	Responsive	20.80%	59.5%	60.0%* Worse than national aggregate	●
89 Cases	Percentage of patients receiving oxygen in which this was prescribed to a stipulated target oxygen saturation (SpO2) range (of 88-92% or 94-98%)	Effective	96.60%	99.3%	100.0%† Worse than national aggregate	●
15 Cases	Percentage of patients receiving non invasive ventilation (NIV) within the first 24 hours of arrival who do so within 3 hours of arrival	Effective	Insufficient data	23.1%	31.9%† Insufficient data	■
52 Cases	Percentage of documented current smokers prescribed smoking-cessation pharmacotherapy	Responsive	57.70%	50.0%	63.6%† Better than national aggregate	▲
125 Cases	Percentage of patients for whom a British Thoracic Society, or equivalent, discharge bundle was completed for the admission	Responsive	22.40%	77.1%	60.0%* Worse than national aggregate	●
125 Cases	Percentage of patients with spirometry confirming FEV1/FVC ratio <0.7 recorded in case file	Effective	31.20%	44.4%	60.3%† Worse than national aggregate	●



[www.rcplondon.ac.uk/projects/national-copd-audit-programme-2013-18](http://www.rcplondon.ac.uk/projects/national-copd-audit-programme-2013-18)

Produced by HQIP  
in partnership with the

